Table 1.A. - Preliminary Cod Landings by Stock Area

| | | | | | | | | | 2003/20 | 004* | 2002/2 | 003* | Fishing |
|------------------------|----------|---------|----------|---------|----------|---------|----------|---------|----------------|----------|-------------|----------|----------|
| | AUG - | - 2003 | SEP - | 2003 | OCT - | 2003 | MAY 03 - | OCT 03 | May03-Oct03 as | Target | May02-Oct02 | Target | Year* |
| | | | | | | | | | a % of Target | TAC | as a % of | TAC | Landings |
| | 1000 Lbs | Percent | TAC | 1000 Lbs | Target TAC | 1000 Lbs | 1000 Lbs |
| GULF OF MAINE | 349 | 52% | 456 | 43% | 661 | 50% | 3,364 | 37% | 57% | 5,899 | 79% | 4,229 | |
| OTTER TRAWL | 165 | 24% | 234 | 22% | 326 | 25% | 1,803 | 20% | 31% | | 37% | | |
| GILLNET | 180 | 27% | 204 | 19% | 318 | 24% | 1,479 | 16% | 25% | | 39% | | |
| HOOK | 0 | 0% | 10 | 1% | 8 | 1% | 31 | 0% | | | 1% | | |
| OTHER GEARS | 4 | 1% | 8 | 1% | 9 | 1% | 51 | 1% | 1% | | 2% | | |
| | | | | | | | | | | | | | |
| GEORGES BANK | 324 | 48% | 598 | 57% | 661 | 50% | 5,806 | 63% | 102% | 5,694 | 201% | 10,803 | |
| OTTER TRAWL | 229 | | 382 | 36% | 585 | 44% | 4,434 | | | | 145% | | |
| GILLNET | 87 | 13% | 125 | 12% | 51 | 4% | 1,089 | 12% | | | 42% | | |
| HOOK | 5 | | 68 | 6% | 13 | 1% | 119 | | | | 8% | | |
| OTHER GEARS | 3 | 0% | 23 | 2% | 14 | 1% | 164 | 2% | 3% | | 6% | | |
| | | T | | | T | | | | | | 1 | | |
| ALL AREAS | 674 | 100% | 1,055 | 100% | 1,323 | 100% | 9,170 | 100% | 79% | 11,593 | 279% | 15,032 | |
| OTTER TRAWL | 394 | | 616 | 58% | 911 | 69% | 6,237 | 68% | | | 182% | | |
| GILLNET | 267 | 40% | 329 | 31% | 369 | 28% | 2,569 | | | | 81% | | |
| HOOK | 5 | | 79 | 7% | 20 | 2% | 150 | 2% | | | 9% | | |
| OTHER GEARS | 7 | 1% | 31 | 3% | 22 | 2% | 215 | 2% | 2% | | 8% | | |
| LANDINGS - ALL AREAS | | | | | | | | | | | | | |
| Fishing Year 2003/2004 | 674 | | 1,055 | | 1,323 | | 9,170 | | | | | | - |
| Fishing Year 2002/2003 | 1,810 | | 1,333 | | 1,229 | | 11,801 | | | | | | 25,327 |
| Fishing Year 2001/2002 | 2,335 | | 2,006 | | 2,000 | | 16,815 | | | | | | 35,510 |
| Fishing Year 2000/2001 | 1,828 | | 1,360 | | 1,530 | | 13,349 | | | | | | 27,945 |
| Fishing Year 1999/2000 | 1,609 | | 1,032 | | 853 | | 11,331 | | | | | | 21,323 |
| Fishing Year 1998/1999 | 1,356 | | 1,486 | | 1,102 | | 12,220 | | | | | | 23,702 |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Cod stock areas: Gulf of Maine: 464-465, 467, 511-515

Georges Bank: 521-637

- 2. Landings in live weight.
- 3. State landings for 2002 have been updated and are complete.
- 4. State landings for CT are estimated for the MAY 2003 OCT 2003 period. State landings for NY are estimated for the JUN 2003 period.
- 5. Gear data are based on vessel trip reports.
- * Fishing Year is May 1 through April 30. The 2003/2004 Fishing Year Target TAC is the same as the 2002/2003 Target TAC because no 2003/2004 Target TAC numbers are available.

Table 1.B. - Preliminary Cod Landings by Major Ports and States, May - Oct 2003

| | | | STOCK | AREAS | | | | | GEAR | TYPES | | | |
|------------------|--------------|----------|---------|----------|---------|----------|---------|----------|---------|----------|---------|----------|---------|
| PORT/ STATE | MAY - OCT 03 | GULF OF | MAINE | GEORGE | S BANK | OTTER | TRAWL | GILL | NET | но | ок | OTHER | GEARS |
| | 1000 Lbs | 1000 Lbs | Percent |
| Portland, ME | 654 | 436 | 67% | 218 | 33% | 365 | 56% | 279 | 43% | 10 | 1% | 0 | 0% |
| Gloucester, MA | 1,993 | 1,314 | 66% | 678 | 34% | 1,387 | 70% | 570 | 29% | 21 | 1% | 14 | 1% |
| Boston, MA | 180 | 24 | 14% | 155 | 86% | 180 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| New Bedford, MA | 3,167 | 77 | 2% | 3,090 | 98% | 3,162 | 100% | 3 | 0% | 0 | 0% | 2 | 0% |
| Point Judith, RI | 33 | 1 | 4% | 32 | 96% | 30 | 90% | 1 | 0% | 0 | 0% | 3 | 8% |
| | | | | | | | | | | | | | |
| MAINE | 1,024 | 805 | 79% | 219 | 21% | 565 | 55% | 444 | 43% | 10 | 1% | 5 | 0% |
| NEW HAMPSHIRE | 663 | 663 | 100% | 0 | 0% | 214 | 32% | 443 | 67% | 0 | 0% | 6 | 1% |
| MASSACHUSETTS | 7,339 | 1,894 | 26% | 5,445 | 74% | 5,360 | 73% | 1,680 | 23% | 140 | 2% | 159 | 2% |
| RHODE ISLAND | 135 | 1 | 1% | 134 | 99% | 92 | 68% | 2 | 1% | 0 | 0% | 42 | 31% |
| CONNECTICUT | 6 | 0 | 0% | 6 | 100% | 6 | 99% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW YORK | 3 | 0 | 0% | 3 | 100% | 0 | 0% | 0 | 0% | 0 | 0% | 3 | 98% |
| NEW JERSEY | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| OTHER NORTHEAST | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| TOTAL | 9,170 | 3,364 | 37% | 5,806 | 63% | 6,237 | 68% | 2,569 | 28% | 150 | 2% | 215 | 2% |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Cod stock areas: Gulf of Maine: 464-465, 467, 511-515

Georges Bank: 521-637

2. State landings for CT are estimated for the May 2003 - Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.

- 3. Landings in live weight.
- 4. Gear data are based on vessel trip reports.

Table 2.A. - Preliminary Haddock Landings by Stock Area

| | | | | | | | | | 2003/20 | 004* | 2002/20 | 003* | Fishing |
|------------------------|----------|---------|----------|---------|----------|---------|----------|---------|----------------|----------|----------------|----------|----------|
| | AUG - | 2003 | SEP - | 2003 | OCT · | 2003 | MAY 03 - | OCT 03 | MAY03-OCT03 | Target | MAY02-OCT02 | Target | Year* |
| | | | | | | | | | as % of Target | TAC | as % of Target | TAC | Landings |
| | 1000 Lbs | Percent | TAC | 1000 Lbs | TAC | 1000 Lbs | 1000 Lbs |
| GULF OF MAINE | 125 | 28% | 123 | 20% | 143 | 15% | 883 | 14% | | | | | |
| OTTER TRAWL | 77 | 17% | 111 | 19% | 120 | 13% | 746 | 12% | | | | | |
| GILLNET | 45 | 10% | 2 | 0% | 17 | 2% | 105 | 2% | | | | | |
| HOOK | 0 | 0% | 9 | 1% | 6 | 1% | 29 | 0% | | | | | |
| OTHER GEARS | 2 | 0% | 0 | 0% | 0 | 0% | 4 | 0% | | | | | |
| | | | | | | | | | | | | | |
| GEORGES BANK | 323 | 72% | 478 | 80% | 805 | 85% | 5,312 | 86% | 21% | 25,586 | 31% | 25,751 | |
| OTTER TRAWL | 319 | 71% | 419 | 70% | 683 | 72% | 5,030 | 81% | 20% | | 29% | | |
| GILLNET | 3 | 1% | 2 | 0% | 1 | 0% | 68 | 1% | 0% | | 1% | | |
| HOOK | 1 | 0% | 58 | 10% | 121 | 13% | 183 | 3% | 1% | | 0% | | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 30 | 0% | 0% | | 0% | | |
| | | | | | | | | | | | | | |
| SNE/ MID-ATLANTIC | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | | | |
| OTTER TRAWL | 0 | 0% | 0 | 0% | 0 | | 0 | 0% | | | | | |
| GILLNET | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | | | |
| НООК | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | | | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | | | |
| _ | | | | | | | | | | | ļ | | |
| ALL AREAS | 448 | 100% | 601 | 100% | 949 | 100% | 6,195 | 100% | | | | | |
| OTTER TRAWL | 396 | 88% | 530 | 88% | 802 | 85% | 5,776 | 93% | | | | | |
| GILLNET | 48 | 11% | 4 | 1% | 19 | | 173 | 3% | | | | | |
| HOOK | 1 | 0% | 67 | 11% | 127 | 13% | 212 | 3% | | | | | |
| OTHER GEARS | 2 | 0% | 0 | 0% | 0 | 0% | 34 | 1% | | | | | |
| LANDINGS - ALL AREAS | | | | | | | | | | | | | |
| Fishing Year 2003/2004 | 448 | | 601 | | 949 | | 6,195 | | | | | | - |
| Fishing Year 2002/2003 | 960 | | 1,149 | | 1,317 | | 8,422 | | | | | | 16,198 |
| Fishing Year 2001/2002 | 1,004 | | 1,205 | | 1,151 | | 5,742 | | | | | | 14,341 |
| Fishing Year 2000/2001 | 683 | | 399 | | 874 | | 3,828 | | | | | | 10,501 |
| Fishing Year 1999/2000 | 622 | | 617 | | 628 | | 4,050 | | | | | | 8,999 |
| Fishing Year 1998/1999 | 561 | | 739 | | 561 | | 3,282 | | | | | | 6,009 |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Haddock stock areas: Gulf of Maine: 464-465, 467, 511-515

Georges Bank: 521-526, 533-534, 537-539, 541-543, 551-552, 561-562

Southern New England / Mid-Atlantic: 611-639

- 2. Landings in live weight.
- 3. State landings for 2002 have been updated and are complete.
- 4. State landings for CT are estimated for the MAY 2003 OCT 2003 period. State landings for NY are estimated for the JUN 2003 period.
- 5. Gear data are based on vessel trip reports.
- * Fishing Year is May 1 through April 30. The 2003/2004 Fishing Year Target TAC is the same as the 2002/2003 Target TAC because no 2003/2004 Target TAC numbers are available.

Table 2.B. - Preliminary Haddock Landings by Major Ports and States, May - Oct 2003

| | | STOCK AREAS | | | | | | | | | GEAR | TYPES | | | |
|------------------|--------------|-------------|---------|----------|---------|----------|---------|----------|---------|----------|---------|----------|---------|----------|---------|
| PORT/ STATE | MAY - OCT 03 | GULF OF | MAINE | GEORGE | S BANK | MID-AT | LANTIC | OTTER | TRAWL | GILL | NET | нс | ок | OTHER | GEARS |
| | 1000 Lbs | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent |
| Portland, ME | 428 | 165 | 38% | 263 | 62% | 0 | 0% | 372 | 87% | 41 | 10% | 14 | 3% | 0 | 0% |
| Gloucester, MA | 1,598 | 397 | 25% | 1,200 | 75% | 0 | 0% | 1,546 | 97% | 35 | 2% | 15 | 1% | 3 | 0% |
| Boston, MA | 144 | 59 | 41% | 85 | 59% | 0 | 0% | 144 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| New Bedford, MA | 3,210 | 55 | 2% | 3,155 | 98% | 0 | 0% | 3,210 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| Point Judith, RI | 83 | 0 | 0% | 83 | 100% | 0 | 0% | 83 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| | | | | | | | | | | | | | | | |
| MAINE | 480 | 217 | 45% | 263 | 55% | 0 | 0% | 422 | 88% | 44 | 9% | 14 | 3% | 0 | 0% |
| NEW HAMPSHIRE | 57 | 57 | 100% | 0 | 0% | 0 | 0% | 30 | 53% | 27 | 47% | 0 | 0% | 0 | 0% |
| MASSACHUSETTS | 5,421 | 608 | 11% | 4,813 | 89% | 0 | 0% | 5,117 | 94% | 102 | 2% | 198 | 4% | 4 | 0% |
| RHODE ISLAND | 224 | 0 | 0% | 224 | 100% | 0 | 0% | 195 | 87% | 0 | 0% | 0 | 0% | 30 | 13% |
| CONNECTICUT | 12 | 0 | 0% | 12 | 100% | 0 | 0% | 12 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW YORK | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW JERSEY | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| OTHER NORTHEAST | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| TOTAL | 6,195 | 883 | 14% | 5,312 | 86% | 0 | 0% | 5,776 | 93% | 173 | 3% | 212 | 3% | 34 | 1% |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Haddock stock areas: Gulf of Maine: 464-465, 467, 511-515

Georges Bank: 521-526, 533-534, 537-539, 541-543, 551-552, 561-562

Southern New England / Mid-Atlantic: 611-639

2. State landings for CT are estimated for the May 2003 - Oct 2003 period. State landings for NY are estimated for the Jun. 2003 period.

- 3. Landings in live weight.
- 4. Gear data are based on vessel trip reports.

Table 3.A. - Preliminary Yellowtail Landings by Stock Area

| | | | | | | | | | 2003/20 | 004* | 2002/20 | 003* | Fishing |
|------------------------|----------|---------|----------|---------|----------|---------|----------|---------|----------------|----------|----------------|----------|----------|
| | AUG - | 2003 | SEP - | 2003 | OCT - 2 | 2003 | MAY 03 - | OCT 03 | MAY03-OCT03 | Target | MAY02-OCT02 | Target | Year* |
| | | | | | | | | | as % of Target | TAC | as % of Target | TAC | Landings |
| | 1000 Lbs | Percent | TAC | 1000 Lbs | TAC | 1000 Lbs | 1000 Lbs |
| GULF OF MAINE | 0 | 0% | 3 | 1% | 5 | 1% | 69 | 2% | | | | | |
| OTTER TRAWL | 0 | 0% | 3 | 1% | 5 | 1% | 69 | 2% | | | | | |
| GILLNET | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | | | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | | | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | | | |
| | | | | | | | | | | | | | |
| CAPE COD | 153 | 39% | 163 | 56% | 324 | 64% | 1,405 | 34% | | | | | |
| OTTER TRAWL | 144 | 37% | 162 | 56% | 324 | 64% | 1,374 | 34% | | | | | |
| GILLNET | 9 | 2% | 1 | 0% | 0 | 0% | 29 | 1% | | | | | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | | | |
| OTHER GEARS | 0 | 0% | 1 | 0% | 0 | 0% | 1 | 0% | | | | | |
| | | | | | | | | | | | | | |
| GEORGES BANK | 236 | 60% | 92 | 32% | 158 | 31% | 2,452 | 60% | 14% | 17,288 | 14% | 15,003 | |
| OTTER TRAWL | 236 | 60% | 92 | 32% | 158 | 31% | 2,448 | 60% | | | 14% | | |
| GILLNET | 0 | 0% | 0 | 0% | 0 | 0% | 2 | 0% | 0% | | 0% | | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0% | | 0% | | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 2 | 0% | 0% | | 0% | | |
| | | | | | | | | | | | ļ | | |
| SNE/ MID-ATLANTIC | 3 | 1% | 32 | 11% | 22 | 4% | 146 | 4% | 37% | 393 | 2% | 2,092 | |
| OTTER TRAWL | 3 | 0 | 32 | 0 | 22 | 0 | 145 | 4% | | | 2% | | |
| GILLNET | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0% | | 0% | | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0% | | 0% | | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 1 | 0% | 0% | | 0% | | |
| | | | | | | | | | | | | | |
| ALL AREAS | 392 | 100% | 291 | 100% | 509 | 100% | 4,072 | 100% | | | | | |
| OTTER TRAWL | 383 | 98% | 289 | 99% | 509 | 100% | 4,035 | 99% | | | | | |
| GILLNET | 9 | 2% | 1 | 0% | 0 | 0% | 32 | 1% | | | | | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | | | |
| OTHER GEARS | 0 | 0% | 1 | 0% | 0 | 0% | 4 | 0% | | | | | |
| LANDINGS - ALL AREAS | | | 204 | | | | 4.070 | | | | | | |
| Fishing Year 2003/2004 | 392 | | 291 | | 509 | | 4,072 | | | | | | |
| Fishing Year 2002/2003 | 485 | | 681 | | 710 | | 4,967 | | | | | | 11,774 |
| Fishing Year 2001/2002 | 853 | | 583 | | 913 | | 7,010 | | | | | | 14,604 |
| Fishing Year 2000/2001 | 547 | | 535 | | 542 | | 5,545 | | | | | | 15,990 |
| Fishing Year 1999/2000 | 281 | | 225 | | 328 | | 3,139 | | | | | | 11,601 |
| Fishing Year 1998/1999 | 410 | | 598 | | 1,184 | | 3,750 | | | | | | 9,571 |

The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Yellowtail stock areas: Gulf of Maine: 464-465, 467, 511-513, 515

Cape Cod: 514, 521

Georges Bank: 522, 525, 542-543, 551-552, 561-562

Southern New England / Mid-Atlantic: 526, 533-534, 537-539, 541, 611-639

- 2. Landings in live weight.
- 3. State landings for 2002 have been updated and are complete.
- 4. State landings for CT are estimated for the MAY 2003 OCT 2003 period. State landings for NY are estimated for the JUN 2003 period.
- 5. Gear data are based on vessel trip reports.
- Fishing Year is May 1 through April 30. The 2003/2004 Fishing Year Target TAC is the same as the 2002/2003 Target TAC because no 2003/2004 Target TAC numbers are available.

Table 3.B. - Preliminary Yellowtail Landings by Major Ports and States, May - Oct 2003

| | | | | | STOCK | AREAS | | | | | | | GEAR | TYPES | | | |
|------------------|--------------|----------|---------|----------|---------|----------|---------|----------|----------------|----------|---------|----------|---------|----------|---------|----------|---------|
| PORT/ STATE | MAY - OCT 03 | GULF OF | MAINE | CAPE | COD | GEORGE | S BANK | | RN NEW LAND | OTTER | TRAWL | GILL | NET | НС | ок | OTHER | GEARS |
| | 1000 Lbs | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent |
| Portland, ME | 36 | 7 | 21% | 3 | 9% | 25 | 70% | 0 | 0% | 36 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| Gloucester, MA | 429 | 11 | 3% | 338 | 79% | 79 | 19% | 1 | 0% | 416 | 97% | 13 | 3% | 0 | 0% | 0 | 0% |
| Boston, MA | 12 | 2 | 14% | 1 | 0% | 9 | 79% | 0 | 0% | 12 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| New Bedford, MA | 2,872 | 3 | 0% | 611 | 21% | 2,204 | 77% | 54 | 2% | 2,871 | 100% | 0 | 0% | 0 | 0% | 1 | 0% |
| Point Judith, RI | 124 | 18 | 14% | 5 | 4% | 52 | 42% | 49 | 40% | 124 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| | | | | | | | | | | | | | | | | | |
| MAINE | 36 | 8 | 22% | 3 | 9% | 25 | 69% | 0 | 0% | 36 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW HAMPSHIRE | 43 | 27 | 63% | 16 | 37% | 0 | 0% | 0 | 0% | 42 | 99% | 0 | 0% | 0 | 0% | 0 | 0% |
| MASSACHUSETTS | 3,756 | 16 | 0% | 1,381 | 37% | 2,300 | 61% | 59 | 2% | 3,722 | 99% | 32 | 1% | 0 | 0% | 2 | 0% |
| RHODE ISLAND | 214 | 18 | 8% | 5 | 2% | 123 | 57% | 68 | 32% | 213 | 99% | 0 | 0% | 0 | 0% | 1 | 1% |
| CONNECTICUT | 5 | 0 | 0% | 0 | 0% | 4 | 93% | 0 | 0% | 4 | 93% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW YORK | 16 | 0 | 0% | 0 | 0% | 0 | 0% | 16 | 100% | 16 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW JERSEY | 2 | 0 | 0% | 0 | 0% | 0 | 0% | 2 | 100% | 2 | 94% | 0 | 0% | 0 | 0% | 0 | 0% |
| OTHER NORTHEAST | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| TOTAL | 4,072 | 69 | 2% | 1,405 | 34% | 2,452 | 60% | 146 | 4% | 4,035 | 99% | 32 | 1% | 0 | 0% | 4 | 0% |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Yellowtail stock areas: Gulf of Maine: 464-465, 467, 511-513, 515

Cape Cod: 514, 521

Georges Bank: 522, 525, 542-543, 551-552, 561-562

Southern New England / Mid-Atlantic: 526, 533-534, 537-539, 541, 611-639

- 2. State landings for CT are estimated for the May 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 3. Landings in live weight.
- 4. Gear data are based on vessel trip reports.

Table 4.A. - Preliminary Prorated White Hake Landings

| | AUG - | 2003 | SEP - | 2003 | ост - | 2003 | MAY 03 - | OCT 03 | Fishing Year* Landings |
|------------------------|------------|---------|------------|-----------|------------|-----------|----------------|-----------|------------------------------|
| | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs |
| SNE/ NORTH | 660 | 100% | 723 | 100% | 789 | 100% | 4,619 | 100% | |
| OTTER TRAWL | 91 | 14% | 407 | 56% | 511 | 65% | 2,565 | 56% | |
| GILLNET | 566 | 86% | 269 | 37% | 276 | 35% | 1,997 | 43% | |
| HOOK | 0 | 0% | 46 | 6% | 2 | 0% | 52 | 1% | |
| OTHER GEARS | 2 | 0% | 1 | 0% | 0 | 0% | 5 | 0% | |
| MID ATLANTIC | • | 00/ | | 00/ | | 00/ | 0 1 | 00/ | |
| MID-ATLANTIC | 0 | 0% | 0 | 0% | 0 | 0% | 2 | 0% | |
| OTTER TRAWL | 0 | 0% | 0 | 0% | 0 | 0% | 1 | 0% | |
| GILLNET | 0 | 0% | 0 | 0% | 0 | 0% | 1 | 0% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| ALL AREAS | 660 | 100% | 723 | 100% | 789 | 100% | 4,620 | 100% | |
| OTTER TRAWL | 91 | 14% | 407 | 56% | 511 | 65% | | 56% | |
| GILLNET | 566 | 86% | 269 | 37% | 276 | 35% | 2,566 1,997 | 43% | |
| HOOK | | 0% | 269 46 | 37% 6% | | 35% 0% | 1,997 | 43% 1% | |
| OTHER GEARS | 0 | 0% | 40 | 0% | 2 | 0% | 52 | 0% | |
| LANDINGS - ALL AREAS | ۷ | 0% | ı | 0% | U | 0% | ວ | 070 | |
| | 660 | | 723 | | 789 | | 4 620 | | |
| Fishing Year 2003/2004 | 502 | | 685 | | 709 706 | | 4,620 | | 7 696 |
| Fishing Year 2002/2003 | 681 | | 972 | | 822 | | 3,159 | | 7,686 |
| Fishing Year 2001/2002 | 652 | | 972 818 | | 782 | | 4,298 | | 8,344 |
| Fishing Year 2000/2001 | | | | | _ | | 4,200 | | 6,932 5,039 |
| Fishing Year 1999/2000 | 762 237 | | 822 222 | | 722 567 | | 3,857 1,773 | | 5,928 5,655 |
| Fishing Year 1998/1999 | 231 | | ZZZ | | 700 | | 1,773 | | 5,655 |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

White Hake Stock Areas: Southern New England/ North: 464-465, 467, 500-562

Mid-Atlantic: 600-639

- 2. Landings in live weight.
- 3. State landings for CT are estimated for the May. 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 5. Gear data are based on vessel trip reports.
- * Fishing Year is May 1 through April 30.

Table 4.B. - Preliminary White Hake Landings by Major Ports and States, May - Oct 2003

| | | | STOCK AREAS SOUTHERN NEW | | | | | | GEAR | TYPES | | | |
|------------------|--------------|----------------|--------------------------|----------|---------|----------|---------|----------|---------|----------|---------|----------|---------|
| PORT/ STATE | MAY - OCT 03 | SOUTHE ENGL | | MID-ATI | LANTIC | OTTER | TRAWL | GILL | NET | но | ок | OTHER | GEARS |
| | 1000 Lbs | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent |
| Portland, ME | 2,335 | 2,335 | 100% | 1 | 0% | 1,478 | 63% | 827 | 35% | 30 | 1% | 0 | 0% |
| Gloucester, MA | 1,108 | 1,108 | 100% | 0 | 0% | 501 | 45% | 597 | 54% | 7 | 1% | 3 | 0% |
| Boston, MA | 177 | 177 | 100% | 0 | 0% | 177 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| New Bedford, MA | 93 | 93 | 100% | 0 | 0% | 93 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| Point Judith, RI | 2 | 2 | 100% | 0 | 0% | 2 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| | | | | | | | | | | | | | |
| MAINE | 2,735 | 2,735 | 100% | 1 | 0% | 1,696 | 62% | 1,008 | 37% | 30 | 1% | 1 | 0% |
| NEW HAMPSHIRE | 411 | 411 | 100% | 0 | 0% | 55 | 13% | 356 | 87% | 0 | 0% | 0 | 0% |
| MASSACHUSETTS | 1,444 | 1,444 | 100% | 0 | 0% | 787 | 54% | 633 | 44% | 21 | 1% | 3 | 0% |
| RHODE ISLAND | 28 | 28 | 100% | 0 | 0% | 27 | 95% | 0 | 0% | 0 | 0% | 1 | 5% |
| CONNECTICUT | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW YORK | 1 | 0 | 0% | 1 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW JERSEY | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| OTHER NORTHEAST | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| TOTAL | 4,620 | 4,619 | 100% | 2 | 0% | 2,566 | 56% | 1,997 | 43% | 52 | 1% | 5 | 0% |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

White Hake stock areas: Southern New England / North: 464-465, 467, 500-562

Mid-Atlantic: 600-639

- 2. State landings for CT are estimated for the May 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 3. Landings in live weight.
- 4. Gear data are based on vessel trip reports.

Table 5.A. - Preliminary Prorated American Plaice Landings

| | AUG - | 2003 | SEP - | 2003 | ост - | 2003 | MAY 03 - | OCT 03 | Fishing Year* Landings |
|------------------------|----------|---------|----------|---------|----------|---------|----------|---------|------------------------------|
| | 1000 Lbs | Percent | 1000 Lbs |
| GULF OF MAINE/ GB | 363 | 100% | 540 | 100% | 438 | 100% | 2,833 | 99% | |
| OTTER TRAWL | 360 | 99% | 537 | 99% | 438 | 100% | 2,812 | 99% | |
| GILLNET | 2 | 1% | 0 | 0% | 0 | 0% | 12 | 0% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 0 | 0% | 3 | 1% | 0 | 0% | 9 | 0% | |
| | | | | | | | | | |
| SNE / MID-ATLANTIC | 0 | 0% | 2 | 0% | 0 | 0% | 18 | 1% | |
| OTTER TRAWL | 0 | 0% | 2 | 0% | 0 | 0% | 18 | 1% | |
| GILLNET | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| | | | • | | | | | | |
| ALL AREAS | 363 | 100% | 542 | 100% | 438 | 100% | 2,851 | 100% | |
| OTTER TRAWL | 360 | 99% | 538 | 99% | 438 | 100% | 2,830 | 99% | |
| GILLNET | 2 | 1% | 0 | 0% | 0 | 0% | 12 | 0% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 0 | 0% | 3 | 1% | 0 | 0% | 9 | 0% | |
| LANDINGS - ALL AREAS | | | | | | | | | |
| Fishing Year 2003/2004 | 363 | | 542 | | 438 | | 2,851 | | - |
| Fishing Year 2002/2003 | 665 | | 757 | | 682 | | 3,711 | | 6,236 |
| Fishing Year 2001/2002 | 1,514 | | 1,226 | | 798 | | 6,807 | | 9,978 |
| Fishing Year 2000/2001 | 1,344 | | 1,311 | | 975 | | 6,490 | | 9,949 |
| Fishing Year 1999/2000 | 1,099 | | 740 | | 580 | | 5,122 | | 7,003 |
| Fishing Year 1998/1999 | 972 | | 722 | | 751 | | 6,263 | | 7,787 |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

American Plaice Stock Areas: Gulf of Maine/ Georges Bank: 464-465, 467, 511-526, 551-552, 561-562 Southern New England/ Mid-Atlantic: 533-534, 537-539, 541-543, 611-639

- 2. Landings in live weight.
- 3. State landings for CT are estimated for the May. 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 4. Gear data are based on vessel trip reports.
- * Fishing Year is May 1 through April 30.

Table 5.B. - Preliminary American Plaice Landings by Major Ports and States, May - Oct 2003

| | | | STOCK | AREAS | | | | | GEAR | TYPES | | | |
|------------------|--------------|----------|---------|----------------|---------|----------|---------|----------|---------|----------|---------|----------|---------|
| PORT/ STATE | MAY - OCT 03 | GULF OF | MAINE | SOUTHE ENGL | | OTTER | TRAWL | GILL | .NET | но | ок | OTHER | GEARS |
| | 1000 Lbs | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent |
| Portland, ME | 688 | 687 | 100% | 1 | 0% | 685 | 100% | 3 | 0% | 0 | 0% | 0 | 0% |
| Gloucester, MA | 420 | 420 | 100% | 0 | 0% | 418 | 100% | 2 | 0% | 0 | 0% | 0 | 0% |
| Boston, MA | 141 | 141 | 100% | 0 | 0% | 141 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| New Bedford, MA | 929 | 914 | 98% | 15 | 2% | 929 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| Point Judith, RI | 32 | 32 | 100% | 0 | 0% | 32 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| | | | | | | | | | | | | | |
| MAINE | 1,067 | 1,066 | 100% | 1 | 0% | 1,060 | 99% | 3 | 0% | 0 | 0% | 3 | 0% |
| NEW HAMPSHIRE | 33 | 33 | 100% | 0 | 0% | 28 | 84% | 5 | 16% | 0 | 0% | 0 | 0% |
| MASSACHUSETTS | 1,675 | 1,660 | 99% | 15 | 1% | 1,672 | 100% | 3 | 0% | 0 | 0% | 0 | 0% |
| RHODE ISLAND | 75 | 73 | 98% | 2 | 2% | 69 | 92% | 0 | 0% | 0 | 0% | 6 | 8% |
| CONNECTICUT | 1 | 1 | 0% | 0 | 0% | 1 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW YORK | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW JERSEY | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| OTHER NORTHEAST | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| TOTAL | 2,851 | 2,833 | 99% | 18 | 1% | 2,830 | 99% | 12 | 0% | 0 | 0% | 9 | 0% |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

American Plaice stock areas: Gulf of Maine/ Georges Bank: 464-465, 467, 511-526, 551-552, 561-562 Southern New England/ Mid-Atlantic: 533-534, 537-539, 541-543, 611-639

- 2. State landings for CT are estimated for the May 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 3. Landings in live weight.
- 4. Gear data are based on vessel trip reports.

Table 6.A. - Preliminary Prorated Winter Flounder Landings

| | AUG - | 2003 | SEP - | 2003 | ост - | 2003 | MAY 03 · | - OCT 03 | Fishing Year* Landings |
|------------------------|-----------|-----------|-----------|-----------|----------|------------|----------|-----------|------------------------------|
| | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs |
| GULF OF MAINE | 117 | 13% | 176 | 19% | 37 | 3% | 704 | 8% | |
| OTTER TRAWL | 98 | 11% | 163 | 17% | 32 | 3% | 624 | 7% | |
| GILLNET | 19 | 2% | 11 | 1% | 5 | 0% | 77 | 1% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 0 | 0% | 3 | 0% | 0 | 0% | 3 | 0% | |
| | | | | | | | | | |
| GEORGES BANK | 480 | 54% | 263 | 28% | 169 | 16% | 5,016 | 59% | |
| OTTER TRAWL | 480 | 54% | 263 | 28% | 169 | 16% | 5,008 | 59% | |
| GILLNET | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 8 | 0% | |
| | | | | | | | | | |
| SNE/ MID-ATLANTIC | 287 | 32% | 506 | 54% | 870 | 81% | 2,715 | 32% | |
| OTTER TRAWL | 277 | 31% | 504 | 53% | 870 | 81% | 2,673 | 32% | |
| GILLNET | 10 | 1% | 2 | 0% | 0 | 0% | 38 | 0% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 3 | 0% | |
| ALL ADEAC | 004 | 4000/ | 0.45 | 4000/ | 4.070 | 4000/ | 0.405 | 4000/ | |
| ALL AREAS | 884 | 100% | 945 | 100% | 1,076 | 100% | 8,435 | 100% | |
| OTTER TRAWL | 855 29 | 97% 3% | 930 13 | 98% 1% | 1,070 | 100% 0% | 8,305 | 98% 1% | |
| GILLNET HOOK | _ | | | | 5 | 0% 0% | 115 | 0% | |
| OTHER GEARS | 0 | 0% 0% | 0 3 | 0% 0% | 0 | 0% | 0 14 | 0% | |
| LANDINGS - ALL AREAS | U | 070 | 3 | 0 70 | U | 0 70 | 14 | 0 70 | |
| Fishing Year 2003/2004 | 884 | | 945 | | 1,076 | | 8,435 | | _ |
| Fishing Year 2002/2003 | 1,841 | | 1,261 | | 861 | | 9,070 | | 11,941 |
| Fishing Year 2001/2002 | 1,926 | | 1,444 | | 1,574 | | 10,180 | | 15,383 |
| Fishing Year 2000/2001 | 1,451 | | 1,031 | | 1,621 | | 8,572 | | 13,796 |
| Fishing Year 1999/2000 | 1,048 | | 956 | | 1,186 | | 6,597 | | 10,366 |
| Fishing Year 1998/1999 | 1,250 | | 1,368 | | 1,387 | | 7,865 | | 11,283 |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Winter Flounder Stock Areas: Gulf of Maine: 464-465, 467, 511-515

Georges Bank: 522, 525, 542-543, 551-552, 561-562

Southern New England/ Mid-Atlantic: 521, 526, 533-534, 537-539, 541, 611-639

- 2. Landings in live weight.
- 3. State landings for CT are estimated for the May. 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 4. Gear data are based on vessel trip reports.
- Fishing Year is May 1 through April 30.

Table 6.B. - Preliminary Winter Flounder Landings by Major Ports and States, May - Oct 2003

| | | | | STOCK | AREAS | | | | | | GEAR | TYPES | | | |
|------------------|--------------|----------|---------|----------|---------|----------------|---------|----------|---------|----------|---------|----------|---------|----------|---------|
| PORT/ STATE | MAY - OCT 03 | GULF O | FMAINE | GEORGE | S BANK | SOUTHE ENGL | | OTTER | TRAWL | GILL | NET | но | ок | OTHER | GEARS |
| | 1000 Lbs | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent |
| Portland, ME | 70 | 7 | 10% | 50 | 70% | 14 | 20% | 70 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| Gloucester, MA | 388 | 203 | 52% | 144 | 37% | 41 | 11% | 351 | 90% | 37 | 10% | 0 | 0% | 0 | 0% |
| Boston, MA | 32 | 4 | 12% | 25 | 80% | 3 | 8% | 32 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| New Bedford, MA | 6,502 | 62 | 1% | 4,726 | 73% | 1,715 | 26% | 6,500 | 100% | 0 | 0% | 0 | 0% | 2 | 0% |
| Point Judith, RI | 309 | 1 | 0% | 35 | 11% | 273 | 88% | 308 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| | | | | | | | | | | | | | | | |
| MAINE | 71 | 7 | 10% | 50 | 70% | 14 | 20% | 71 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW HAMPSHIRE | 10 | 10 | 100% | 0 | 0% | 0 | 0% | 6 | 62% | 4 | 37% | 0 | 0% | 0 | 0% |
| MASSACHUSETTS | 7,428 | 686 | 9% | 4,897 | 66% | 1,846 | 25% | 7,311 | 98% | 111 | 1% | 0 | 0% | 6 | 0% |
| RHODE ISLAND | 431 | 1 | 0% | 67 | 16% | 363 | 84% | 425 | 99% | 0 | 0% | 0 | 0% | 6 | 1% |
| CONNECTICUT | 30 | 0 | 0% | 2 | 8% | 28 | 92% | 30 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW YORK | 129 | 0 | 0% | 0 | 0% | 129 | 100% | 128 | 99% | 0 | 0% | 0 | 0% | 1 | 1% |
| NEW JERSEY | 336 | 0 | 0% | 0 | 0% | 336 | 100% | 335 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| OTHER NORTHEAST | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| TOTAL | 8,435 | 704 | 8% | 5,016 | 59% | 2,715 | 32% | 8,305 | 98% | 115 | 1% | 0 | 0% | 14 | 0% |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Winter Flounder stock areas: Gulf of Maine: 464-465, 467, 511-515

Georges Bank: 522, 525, 542-543, 551-552, 561-562

Southern New England / Mid-Atlantic: 521, 526, 533-534, 537-539, 541, 611-639

- 2. State landings for CT are estimated for the May 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 3. Landings in live weight.
- 4. Gear data are based on vessel trip reports.

Table 7.A. - Preliminary Prorated Witch Flounder Landings

| | AUG - | 2003 | SEP - | 2003 | ост - | 2003 | MAY 03 - | OCT 03 | Fishing Year* Landings |
|------------------------|----------|---------|----------|---------|----------|---------|----------|---------|------------------------------|
| | 1000 Lbs | Percent | 1000 Lbs |
| GULF OF MAINE/ GB | 432 | 97% | 588 | 98% | 494 | 98% | 3,861 | 98% | |
| OTTER TRAWL | 423 | 95% | 585 | 97% | 493 | 98% | 3,819 | 97% | |
| GILLNET | 8 | 2% | 2 | 0% | 0 | 0% | 36 | 1% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 1 | 0% | 1 | 0% | 0 | 0% | 6 | 0% | |
| | | | | | | | | | |
| SNE/ MID-ATLANTIC | 15 | 3% | 15 | 2% | 8 | 2% | 77 | 2% | |
| OTTER TRAWL | 15 | 3% | 15 | 2% | 8 | 2% | 75 | 2% | |
| GILLNET | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 2 | 0% | |
| | | | | | • | | | | |
| ALL AREAS | 446 | 100% | 602 | 100% | 502 | 100% | 3,938 | 100% | |
| OTTER TRAWL | 437 | 98% | 600 | 100% | 501 | 100% | 3,894 | 99% | |
| GILLNET | 8 | 2% | 2 | 0% | 0 | 0% | 36 | 1% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 1 | 0% | 1 | 0% | 0 | 0% | 8 | 0% | |
| LANDINGS - ALL AREAS | | | | | | | | | |
| Fishing Year 2003/2004 | 446 | | 602 | | 502 | | 3,938 | | - |
| Fishing Year 2002/2003 | 708 | | 942 | | 684 | | 3,764 | | 6,168 |
| Fishing Year 2001/2002 | 911 | | 711 | | 574 | | 4,097 | | 7,246 |
| Fishing Year 2000/2001 | 743 | | 567 | | 527 | | 3,379 | | 5,941 |
| Fishing Year 1999/2000 | 743 | | 468 | | 452 | | 3,330 | | 4,896 |
| Fishing Year 1998/1999 | 546 | | 362 | | 337 | | 2,815 | | 4,048 |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Witch Flounder Stock Areas: Gulf of Maine/ Georges Bank: 464-465, 467, 511-526, 551-552, 561-562

Southern New England/ Mid-Atlantic: 533-534, 537-539, 541-543, 611-639

- 2. Landings in live weight.
- 3. State landings for CT are estimated for the May. 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 4. Gear data are based on vessel trip reports.
- Fishing Year is May 1 through April 30.

Table 7.B. - Preliminary Witch Flounder Landings by Major Ports and States, May - Oct 2003

| | | | STOCK | AREAS | | | | | GEAR | TYPES | | | |
|------------------|--------------|----------|---------|----------------|---------|----------|---------|----------|---------|----------|---------|----------|---------|
| PORT/ STATE | MAY - OCT 03 | GULF OF | MAINE | SOUTHE ENGL | | OTTER | TRAWL | GILL | .NET | но | ок | OTHER | GEARS |
| | 1000 Lbs | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent |
| Portland, ME | 438 | 437 | 100% | 1 | 0% | 437 | 100% | 1 | 0% | 0 | 0% | 0 | 0% |
| Gloucester, MA | 1,184 | 1,183 | 100% | 1 | 0% | 1,162 | 98% | 22 | 2% | 0 | 0% | 0 | 0% |
| Boston, MA | 166 | 166 | 100% | 0 | 0% | 166 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| New Bedford, MA | 1,013 | 996 | 98% | 17 | 2% | 1,012 | 100% | 0 | 0% | 0 | 0% | 1 | 0% |
| Point Judith, RI | 50 | 35 | 69% | 15 | 31% | 50 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| | | | | | | | | | | | | | |
| MAINE | 706 | 705 | 100% | 1 | 0% | 704 | 100% | 1 | 0% | 0 | 0% | 1 | 0% |
| NEW HAMPSHIRE | 97 | 97 | 100% | 0 | 0% | 95 | 98% | 2 | 2% | 0 | 0% | 0 | 0% |
| MASSACHUSETTS | 2,982 | 2,962 | 99% | 20 | 1% | 2,946 | 99% | 33 | 1% | 0 | 0% | 3 | 0% |
| RHODE ISLAND | 112 | 93 | 83% | 19 | 17% | 108 | 96% | 0 | 0% | 0 | 0% | 4 | 4% |
| CONNECTICUT | 4 | 4 | 100% | 0 | 0% | 4 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW YORK | 1 | 0 | 0% | 1 | 100% | 1 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW JERSEY | 36 | 0 | 0% | 36 | 100% | 36 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| OTHER NORTHEAST | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| TOTAL | 3,938 | 3,861 | 98% | 77 | 2% | 3,894 | 99% | 36 | 1% | 0 | 0% | 8 | 0% |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Witch Flounder stock areas: Gulf of Maine/ Georges Bank: 464-465, 467, 511-526, 551-552, 561-562

Southern New England/ Mid-Atlantic: 533-534, 537-539, 541-543, 611-639

- 2. State landings for CT are estimated for the May 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 3. Landings in live weight.
- 4. Gear data are based on vessel trip reports.

Table 8.A. - Preliminary Prorated Windowpane Flounder Landings

| | AUG - | 2003 | SEP - | 2003 | OCT - | 2003 | MAY 03 - | OCT 03 | Fishing Year* Landings |
|--------------------------------------------------|----------|---------|----------|---------|----------|---------|-------------|---------|------------------------------|
| | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs |
| GULF OF MAINE/ GB | 0 | 0% | 0 | 0% | 0 | 0% | 3 | 11% | |
| OTTER TRAWL | 0 | 0% | 0 | 0% | 0 | 0% | 3 | 11% | |
| GILLNET | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| | | | | | | | | | |
| SNE/ MID-ATLANTIC | 1 | 97% | 2 | 99% | 2 | 99% | 25 | 89% | |
| OTTER TRAWL | 1 | 97% | 2 | 99% | 2 | 98% | 25 | 88% | |
| GILLNET | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| | | | | | | | | | |
| ALL AREAS | 1 | 100% | 2 | 100% | 3 | 100% | 28 | 100% | |
| OTTER TRAWL | 1 | 100% | 2 | 100% | 2 | 98% | 28 | 99% | |
| GILLNET | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| LANDINGS - ALL AREAS | 4 | | • | | • | | 00 | | |
| Fishing Year 2003/2004 | 1 | | 2 | | 3 | | 28 50 | | - |
| Fishing Year 2002/2003 | 7 | | 3 | | 2 | | 56 | | 135 |
| Fishing Year 2001/2002 | 6 14 | | 9 12 | | 15 7 | | 61 183 | | 261 431 |
| Fishing Year 2000/2001 Fishing Year 1999/2000 | 14 | | 17 | | - | | 183 88 | | 431 600 |
| Fishing Year 1999/2000 Fishing Year 1998/1999 | 34 | | 107 | | 12 52 | | 331 | | 600 613 |
| 1 1311111y 1 cal 1330/1333 | JŦ | | 101 | | JZ | | 33 I | | 013 |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Windowpane Flounder Stock Areas: Gulf of Maine/ Georges Bank: 464-465, 511-515, 521-522, 525, 542-543, 551-552, 561-562 Southern New England/ Mid-Atlantic: 526, 533-534, 537-539, 611-639

- 2. Landings in live weight.
- 3. State landings for CT are estimated for the May. 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 4. Gear data are based on vessel trip reports.
- * Fishing Year is May 1 through April 30.

Table 8.B. - Preliminary Windowpane Flounder Landings by Major Ports and States, May - Oct 2003

| | | | STOCK | AREAS | | | | | GEAR | TYPES | | | |
|------------------|--------------|----------|---------|----------|----------------|----------|---------|----------|---------|----------|---------|----------|---------|
| PORT/ STATE | MAY - OCT 03 | GULF OF | MAINE | | RN NEW _AND | OTTER | TRAWL | GILL | .NET | но | ок | OTHER | GEARS |
| | 1000 Lbs | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent |
| Portland, ME | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| Gloucester, MA | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| Boston, MA | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| New Bedford, MA | 2 | 2 | 96% | 0 | 0% | 2 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| Point Judith, RI | 5 | 1 | 0% | 4 | 82% | 5 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| | | | | | | | | | | | | | |
| MAINE | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW HAMPSHIRE | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| MASSACHUSETTS | 2 | 2 | 96% | 0 | 0% | 2 | 99% | 0 | 0% | 0 | 0% | 0 | 0% |
| RHODE ISLAND | 7 | 1 | 0% | 6 | 85% | 7 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| CONNECTICUT | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW YORK | 12 | 0 | 0% | 12 | 100% | 12 | 98% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW JERSEY | 8 | 0 | 0% | 8 | 100% | 7 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| OTHER NORTHEAST | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| TOTAL | 28 | 3 | 11% | 25 | 89% | 28 | 99% | 0 | 0% | 0 | 0% | 0 | 0% |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Windowpane Flounder stock areas: Gulf of Maine/ Georges Bank: 464-465, 511-515, 521-522, 525, 542-543, 551-552, 561-562 Southern New England/ Mid-Atlantic: 526, 533-534, 537-539, 611-639

- 2. State landings for CT are estimated for the May 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 3. Landings in live weight.
- 4. Gear data are based on vessel trip reports.

Table 9 - Preliminary Multispecies Landings**

| | | | | | | | | | | 2003/2 | 2004* | 2002/2 | 003* | Fishing |
|-----------------------|------------|----------|---------|----------|---------|----------------|---------|----------|---------|--------------------------|---------------|--------------------------|---------------|--------------------------------------------------|
| | | AUG | - 2003 | SEP - | - 2003 | ост - | 2003 | MAY 03 - | OCT 03 | May03-Oct03 as a % of | Target TAC | May02-Oct02 as a % of | Target TAC | Year* Landings |
| | | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | Target TAC | 1000 Lbs | Target TAC | 1000 Lbs | 1000 Lbs |
| ATLANTIC POLLOCK | 2003/2004 | 437 | 15% | 617 | 17% | 768 | 21% | 3,877 | 16% | 6% | | | | |
| | 2002/2003 | 524 | 12% | 590 | 14% | 561 | 16% | 3,053 | 13% | | | 5% | | |
| | 2001/2002 | 910 | 15% | 824 | 16% | 849 | 18% | 4,413 | 15% | | | | | |
| | 2000/ 2001 | 952 | 18% | 1,003 | 21% | 746 | 16% | 4,915 | 17% | | | | | |
| | 1999/ 2000 | 662 | 15% | 934 | 23% | 652 | 18% | 4,017 | 17% | | | | | |
| | | | | | | | | | | | | | | |
| REDFISH | 2003/2004 | 46 | 2% | 215 | 6% | 85 | 2% | 520 | 2% | 1% | | | | |
| | 2002/2003 | 61 | 1% | 31 | 1% | 37 | 1% | 314 | 1% | | | 1% | | |
| | 2001/2002 | 101 | 2% | 53 | 1% | 58 | 1% | 470 | 2% | | | | | |
| | 2000/ 2001 | 67 | 1% | 50 | 1% | 56 | 1% | 440 | 2% | | | | | |
| | 1999/ 2000 | 98 | 2% | 47 | 1% | 37 | 1% | 418 | 2% | | | | | |
| WHITE HAKE | 2003/2004 | 660 | 23% | 723 | 20% | 789 | 22% | 4,620 | 19% | 8% | | | | |
| WILLE HAVE | 2003/2004 | 502 | 12% | 685 | 16% | 789 706 | 20% | 3,159 | 14% | 076 | | 6% | | |
| | | | 11% | | | 822 | | | 14% | — | | 0% | | |
| | 2001/2002 | 681 | | 972 | 19% | | 18% | 4,298 | | | | | | |
| | 2000/ 2001 | 652 | 12% | 818 | 17% | 782 | 17% | 4,200 | 15% | | | | | |
| | 1999/ 2000 | 762 | 17% | 822 | 21% | 722 | 20% | 3,857 | 16% | 1 | | | | - |
| AMERICAN PLAICE | 2003/2004 | 363 | 13% | 542 | 15% | 438 | 12% | 2,851 | 12% | 5% | | 1 | | |
| | 2002/2003 | 665 | 15% | 757 | 18% | 682 | 19% | 3,711 | 16% | | | 7% | | |
| | 2001/2002 | 1.514 | 25% | 1.226 | 23% | 798 | 17% | 6.807 | 22% | | | . ,, | | |
| | 2000/ 2001 | 1,344 | 26% | 1,311 | 27% | 975 | 21% | 6,490 | 23% | | | | | |
| | 1999/ 2000 | 1,099 | 25% | 740 | 19% | 580 | 16% | 5,122 | 22% | | | 1 | | |
| | | , | | | | | | , | | | | | | |
| WINTER FLOUNDER | 2003/2004 | 884 | 31% | 945 | 26% | 1,076 | 29% | 8,435 | 35% | 14% | | | | |
| | 2002/2003 | 1,841 | 43% | 1,261 | 30% | 861 | 24% | 9,070 | 39% | | | 16% | | |
| | 2001/2002 | 1,926 | 32% | 1,444 | 28% | 1,574 | 34% | 10,180 | 34% | | | | | |
| | 2000/ 2001 | 1,451 | 28% | 1,031 | 22% | 1,621 | 34% | 8,572 | 30% | | | | | |
| | 1999/ 2000 | 1,048 | 24% | 956 | 24% | 1,186 | 33% | 6,597 | 28% | | | | | |
| | | | | | | | | | | | | | | |
| WITCH FLOUNDER | 2003/2004 | 446 | 16% | 602 | 17% | 502 | 14% | 3,938 | 16% | 6% | | | | |
| | 2002/2003 | 708 | 16% | 942 | 22% | 684 | 19% | 3,764 | 16% | | | 7% | | |
| | 2001/2002 | 911 | 15% | 711 | 14% | 574 | 12% | 4,097 | 14% | | | | | |
| | 2000/ 2001 | 743 | 14% | 567 | 12% | 527 | 11% | 3,379 | 12% | | | | | |
| | 1999/ 2000 | 743 | 17% | 468 | 12% | 452 | 12% | 3,330 | 14% | | | | | |
| WINDOWPANE FLDR | 2003/2004 | 1 | 0% | 2 | 0% | 3 | 0% | 28 | 0% | 0% | | | | - |
| THE STATE I LDR | 2003/2004 | 7 | 0% | 3 | 0% | 2 | 0% | 56 | 0% | 0 /0 | | 0% | | 1 |
| | 2002/2003 | 6 | 0% | 9 | 0% | <u>∠</u> 15 | 0% | 61 | 0% | + | | U% | | |
| | | | | | | 7 | 0% | | | + | | 1 | | |
| | 2000/ 2001 | 14 | 0% | 12 | 0% | | | 183 | 1% | + | | | | 1 |
| | 1999/ 2000 | 10 | 0% | 17 | 0% | 12 | 0% | 88 | 0% | | | | | |
| ALL 7 SPECIES - TOTAL | 2003/ 2004 | 2,838 | 100% | 3,648 | 100% | 3,661 | 100% | 24,270 | 100% | 40% | 61,329 | | | - |
| | 2002/ 2003 | 4,309 | 100% | 4,269 | 100% | 3,532 | 100% | 23,128 | 100% | | | 41% | 56,200 | 41,057 |
| | 2001/ 2002 | 6,049 | 100% | 5,240 | 100% | 4,690 | 100% | 30,326 | 100% | | | | | 52,175 |
| | 2000/ 2001 | 5,222 | 100% | 4,792 | 100% | 4,714 | 100% | 28,179 | 100% | | | | | 46,273 |
| | 1999/ 2000 | 4,421 | 100% | 3,984 | 100% | 3,641 | 100% | 23,429 | 100% | | | | | 38,302 |

^{1.} Landings in live weight.

Gear data are based on vessel trip reports.

^{3.} State landings for CT are estimated for the May 2003 - Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.

* Fishing Year is May 1 through April 30. The 2003/2004 Fishing Year Target TAC is the same as the 2002/2003 Target TAC because no 2003/2004 Target TAC numbers are available.

^{**} For the 7 species under the Multispecies FMP other than cod, haddock and yellowtail flounder.

Table 10.A. - Preliminary Silver Hake Landings by Stock Area

| | AUG - | 2003 | SEP - | 2003 | ост - | 2003 | MAY 03 - | OCT 03 | Fishing Year* Landings |
|------------------------|----------|---------|----------|---------|----------|---------|----------|---------|------------------------------|
| | 1000 Lbs | Percent | 1000 Lbs |
| GULF OF MAINE | 812 | 68% | 648 | 25% | 411 | 24% | 2,790 | 29% | |
| OTTER TRAWL | 809 | 67% | 644 | 24% | 403 | 24% | 2,764 | 29% | |
| GILLNET | 1 | 0% | 4 | 0% | 8 | 0% | 16 | 0% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 9 | 0% | |
| OTHER GEARS | 1 | 0% | 0 | 0% | 0 | 0% | 1 | 0% | |
| | _ | | | | | | | | |
| SNE/ MID-ATLANTIC | 388 | 32% | 1,992 | 75% | 1,295 | 76% | 6,725 | 71% | |
| OTTER TRAWL | 388 | 32% | 1,991 | 75% | 1,294 | 76% | 6,673 | 70% | |
| GILLNET | 0 | 0% | 0 | 0% | 0 | 0% | 52 | 1% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| | | | | | | | | | |
| ALL AREAS | 1,199 | 100% | 2,640 | 100% | 1,705 | 100% | 9,515 | 100% | |
| OTTER TRAWL | 1,197 | 100% | 2,636 | 100% | 1,697 | 99% | 9,437 | 99% | |
| GILLNET | 1 | 0% | 4 | 0% | 9 | 1% | 67 | 1% | |
| HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 9 | 0% | |
| OTHER GEARS | 1 | 0% | 0 | 0% | 0 | 0% | 1 | 0% | |
| LANDINGS - ALL AREAS | | | | | | | | | |
| Fishing Year 2003/2002 | 1,199 | | 2,640 | | 1,705 | | 9,515 | | - |
| Fishing Year 2002/2003 | 1,359 | | 1,691 | | 1,685 | | 9,684 | | 15,741 |
| Fishing Year 2001/2002 | 2,748 | | 2,788 | | 3,225 | | 16,874 | | 24,585 |
| Fishing Year 2000/2001 | 2,157 | | 2,599 | | 2,931 | | 13,931 | | 27,359 |
| Fishing Year 1999/2000 | 2,695 | | 2,588 | | 3,015 | | 17,510 | | 29,045 |
| Fishing Year 1998/1999 | 3,353 | | 3,069 | | 2,610 | | 17,925 | | 33,308 |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Silver Hake Stock Areas: Gulf of Maine: 464-465, 511-515, 521-522, 561 Southern New England: 525-552, 611-639

- 2. Landings in live weight.
- 3. State landings for CT are estimated for the May 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 4. Gear data are based on vessel trip reports.
- * Fishing Year is May 1 through April 30.

Table 10.B. - Preliminary Silver Hake Landings by Major Ports and States, May 2003 - Oct 2003

| | | | STOCK | AREAS | | | | | GEAR | TYPES | | | |
|------------------|--------------|----------|---------|----------------|---------|----------|---------|----------|---------|----------|---------|----------|---------|
| PORT/ STATE | MAY - OCT 03 | GULF OF | MAINE | SOUTHE ENGL | | OTTER | TRAWL | GILL | .NET | но | ок | OTHER | GEARS |
| | 1000 Lbs | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent |
| Portland, ME | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| Gloucester, MA | 393 | 393 | 100% | 0 | 0% | 372 | 95% | 11 | 3% | 9 | 2% | 0 | 0% |
| Boston, MA | 1 | 1 | 0% | 0 | 0% | 1 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| New Bedford, MA | 4,012 | 1,587 | 40% | 2,425 | 60% | 4,012 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| Point Judith, RI | 2,759 | 369 | 13% | 2,389 | 87% | 2,758 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| | | | | | | | | | | | | | |
| MAINE | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW HAMPSHIRE | 89 | 89 | 100% | 0 | 0% | 87 | 97% | 2 | 3% | 0 | 0% | 0 | 0% |
| MASSACHUSETTS | 4,562 | 2,134 | 47% | 2,429 | 53% | 4,539 | 99% | 13 | 0% | 9 | 0% | 1 | 0% |
| RHODE ISLAND | 3,015 | 370 | 12% | 2,645 | 88% | 2,964 | 98% | 51 | 2% | 0 | 0% | 0 | 0% |
| CONNECTICUT | 824 | 0 | 0% | 824 | 100% | 824 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW YORK | 992 | 197 | 20% | 795 | 80% | 992 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW JERSEY | 32 | 0 | 0% | 32 | 100% | 32 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| OTHER NORTHEAST | 0 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| TOTAL | 9,515 | 2,790 | 29% | 6,725 | 71% | 9,437 | 99% | 67 | 1% | 9 | 0% | 1 | 0% |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Silver Hake stock areas: Gulf of Maine: 464-465, 511-515, 521-522, 561 Southern New England: 525-552, 611-639

- 2. State landings for CT are estimated for the May 2003 Oct 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 3. Landings in live weight.
- 4. Gear data are based on vessel trip reports.

Table 11.A. - Preliminary Monkfish Landings by Stock Area

| | | | | | | | | | | 2003/2 | 004* | 2002/2 | 2003* | Fishing |
|--------------|-----------------|----------|----------|----------|----------|----------|-----------|----------|-----------|-------------|----------|-------------|----------|----------|
| | | AUG - | 2003 | SEP - | 2003 | OCT - | 2003 | MAY 03 - | OCT 03 | May03-Oct03 | Target | May02-Oct02 | Target | Year* |
| | | | | | | | | | | as a % of | TAC | as a % of | TAC | Landings |
| | | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | Target TAC | 1000 Lbs | Target TAC | 1000 Lbs | 1000 Lbs |
| NORTHERN | | 1,628 | 72% | 2,472 | 82% | 3,267 | 66% | 14,025 | 56% | 36% | 39,039 | 56% | 25,737 | |
| | OTTER TRAWL | 264 | 12% | 1,768 | 59% | 2,589 | 53% | 9,077 | 36% | 23% | | 35% | | |
| | GILLNET | 981 | 43% | 673 | | 654 | 13% | 4,279 | 17% | | | 20% | | |
| | HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 1 | 0% | | | 0% | | |
| | OTHER GEARS | 383 | 17% | 31 | 1% | 23 | 0% | 668 | 3% | 2% | | 1% | | |
| COLITIEDA | | 00.4 | 000/ | 504 | 400/ | 4.040 | 0.40/ | 44.405 | 4.40/ | 500/ | 00 544 | 070/ | 47.400 | |
| SOUTHERN | OTTED TO AVAIL | 634 | 28% | 531 | 18% | 1,648 | 34% | 11,165 | 44% | 50% | 22,511 | 27% | 17,463 | |
| ' | OTTER TRAWL | 153 | 7% | 339 | 11% | 251 | 5% | 1,487 | 6% | | | 3% | | |
| | GILLNET HOOK | 112 | 5% 0% | 85 | 3% 0% | 1,100 | 22% 0% | 7,990 | 32% 0% | | | 18% 0% | | |
| | OTHER GEARS | 0 | 16% | 0 107 | 0% 4% | 297 | 6% | 1 600 | 7% | 7% | | 5% | | |
| <u> </u> | OTHER GEARS | 368 | 10% | 107 | 470 | 291 | 070 | 1,688 | 170 | 1 70 | | 3% | | |
| ALL AREAS | | 2,262 | 100% | 3,003 | 100% | 4,915 | 100% | 25,189 | 100% | 41% | 61,551 | 83% | 43,200 | |
| | OTTER TRAWL | 417 | 18% | 2,107 | 70% | 2,841 | 58% | 10,564 | 42% | | - / | 39% | -, | |
| | GILLNET | 1,093 | 48% | 757 | 25% | 1,754 | 36% | 12,269 | 49% | | | 38% | | |
| | HOOK | 0 | 0% | 0 | 0% | 0 | 0% | 1 | 0% | | | 0% | | |
| | OTHER GEARS | 751 | 33% | 138 | 5% | 320 | 7% | 2,356 | 9% | 4% | | 6% | | |
| LANDINGS - A | ALL AREAS | | | | | | | | | | | | | |
| Fishing | Year 2003/2004 | 2,262 | | 3,003 | | 4,915 | | 25,189 | | | | | | - |
| Fishing | Year 2002/2003 | 4,614 | | 3,284 | | 3,047 | | 22,499 | | | | | | 48,077 |
| Fishing | Year 2001/2002 | 5,415 | | 3,727 | | 3,316 | | 26,874 | | | | | | 57,148 |
| | Year 2000/2001 | 3,935 | | 2,897 | | 2,818 | | 20,830 | | | | | | 43,694 |
| | Year 1999/2000 | 6,405 | | 3,208 | | 3,108 | | 33,756 | | | | | | 52,979 |
| Fishing | Year 1998/1999 | 6,015 | | 4,006 | | 3,323 | | 34,124 | | | | | | 59,487 |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Monkfish Stock Areas: Northern: 464-465, 467, 511-515, 521-522, 561-562 Southern: 525-526, 533-534, 537-539, 541-543, 611-639

- 2. Landings in live weight.
- 3. State landings for CT are estimated for the May 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 4. Gear data are based on vessel trip reports.
- * Fishing Year is May 1 through April 30.

Table 11.B. - Preliminary Monkfish Landings by Major Ports and States, May - Oct 2003

| | | | STOCK | AREAS | | | | | GEAR | TYPES | | | |
|------------------|--------------|----------|---------|----------|---------|----------|---------|----------|---------|----------|---------|----------|---------|
| PORT/ STATE | MAY - OCT 03 | NORT | HERN | SOUT | HERN | OTTER | TRAWL | GILL | NET | но | ок | OTHER | GEARS |
| | 1000 Lbs | 1000 Lbs | Percent |
| Portland, ME | 2,868 | 2,821 | 98% | 46 | 2% | 2,340 | 82% | 528 | 18% | 0 | 0% | 0 | 0% |
| Gloucester, MA | 3,448 | 3,382 | 98% | 65 | 2% | 1,360 | 39% | 2,082 | 60% | 0 | 0% | 5 | 0% |
| Boston, MA | 981 | 981 | 100% | 0 | 0% | 981 | 100% | 0 | 0% | 0 | 0% | 0 | 0% |
| New Bedford, MA | 6,117 | 3,071 | 50% | 3,045 | 50% | 2,991 | 49% | 1,218 | 20% | 0 | 0% | 1,908 | 31% |
| Point Judith, RI | 1,575 | 87 | 6% | 1,488 | 94% | 631 | 40% | 855 | 54% | 0 | 0% | 89 | 6% |
| | | | | | | | | | | | | | |
| MAINE | 3,948 | 3,894 | 99% | 54 | 1% | 3,302 | 84% | 639 | 16% | 0 | 0% | 6 | 0% |
| NEW HAMPSHIRE | 907 | 907 | 100% | 0 | 0% | 53 | 6% | 848 | 93% | 0 | 0% | 6 | 1% |
| MASSACHUSETTS | 12,639 | 8,599 | 68% | 4,040 | 32% | 5,778 | 46% | 4,922 | 39% | 0 | 0% | 1,938 | 15% |
| RHODE ISLAND | 3,117 | 623 | 20% | 2,494 | 80% | 1,223 | 39% | 1,799 | 58% | 0 | 0% | 96 | 3% |
| CONNECTICUT | 283 | 0 | 0% | 283 | 100% | 35 | 12% | 201 | 71% | 0 | 0% | 48 | 17% |
| NEW YORK | 382 | 0 | 0% | 382 | 100% | 52 | 14% | 329 | 86% | 0 | 0% | 1 | 0% |
| NEW JERSEY | 3,069 | 0 | 0% | 3,069 | 100% | 90 | 3% | 2,769 | 90% | 0 | 0% | 210 | 7% |
| OTHER NORTHEAST | 843 | 1 | 0% | 842 | 100% | 31 | 4% | 760 | 90% | 0 | 0% | 51 | 6% |
| TOTAL | 25,189 | 14,025 | 56% | 11,165 | 44% | 10,564 | 42% | 12,269 | 49% | 1 | 0% | 2,356 | 9% |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Monkfish stock areas: Northern: 464-465, 467, 511-515, 521-522, 561-562

Southern: 525-526, 533-534, 537-539, 541-543, 611-639

- 2. State landings for CT are estimated for the May 2003 Oct. 2003 period. State landings for NY are estimated for the Jun. 2003 period.
- 3. Landings in live weight.
- 4. Gear data are based on vessel trip reports.

Table 12.A. - Preliminary Sea Scallop Landings by Stock Area

| | AUG - | 2003 | SEP - | 2003 | ост - | 2003 | MAR - 0 | OCT 03 | Fishing Year* Landings |
|------------------------|------------|-----------|----------------|-----------|----------------|-----------|------------------|-----------|------------------------------|
| | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs |
| GULF OF MAINE | 11 | 0% | 23 | 1% | 18 | 0% | 400 | 1% | |
| OTTER TRAWL | 0 | 0% | 0 | 0% | 0 | 0% | 3 | 0% | |
| DREDGE | 11 | 0% | 22 | 1% | 17 | 0% | 395 | 1% | |
| OTHER GEARS | 0 | 0% | 1 | 0% | 1 | 0% | 2 | 0% | |
| | | | | | | | | | |
| SOUTH CHANNEL | 868 | 15% | 445 | 12% | 238 | 4% | 4,964 | 11% | |
| OTTER TRAWL | 1 | 0% | 1 | 0% | 1 | 0% | 29 | 0% | |
| DREDGE | 868 | 15% | 444 | 12% | 237 | 4% | 4,935 | 11% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| GEORGES BANK NORTH | 374 | 6% | 0 | 0% | 8 | 0% | 2.449 | 5% | |
| OTTER TRAWL | 0 | 0% | 0 | 0% | 0 | 0% | 2,449 | 0% | |
| DREDGE | 374 | 6% | 0 | 0% | 8 | 0% | 2,449 | 5% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 2,443 | 0% | |
| SHIER GEARG | <u> </u> | 0 70 | o _l | 0 70 | o _l | 0 70 | O ₁ | 0 70 | |
| GEORGES BANK SOUTH | 352 | 6% | 92 | 2% | 108 | 2% | 1,309 | 3% | |
| OTTER TRAWL | 0 | 0% | 0 | 0% | 0 | 0% | 1 | 0% | |
| DREDGE | 352 | 6% | 92 | 2% | 108 | 2% | 1,308 | 3% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| | | | | | | | | | |
| S. NEW ENGLAND | 0 | 0% | 0 | 0% | 1 | 0% | 105 | 0% | |
| OTTER TRAWL | 0 | 0% | 0 | 0% | 1 | 0% | 6 | 0% | |
| DREDGE | 0 | 0% | 0 | 0% | 0 | 0% | 99 | 0% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| | | | | | | | | | |
| NEW YORK BIGHT | 1,037 | 18% | 1,526 | 40% | 2,358 | 43% | 16,471 | 36% | |
| OTTER TRAWL | 2 | 0% | 4 500 | 0% | 2 | 0% | 172 | 0% | |
| DREDGE OTHER GEARS | 1,035 | 18% 0% | 1,522 | 40% 0% | 2,356 | 43% | 16,299 | 36% 0% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | U | 0% | 0 | 0% | |
| DE / MD / VA | 3,063 | 53% | 1.729 | 45% | 2.778 | 50% | 19,737 | 43% | |
| OTTER TRAWL | 357 | 6% | 144 | 4% | 278 | 5% | 2,460 | 5% | |
| DREDGE | 2,705 | 46% | 1,585 | 41% | 2,500 | 45% | 17,277 | 38% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| | - 1 | | - | | - | | | | |
| VA / NC | 129 | 2% | 16 | 0% | 28 | 1% | 380 | 1% | |
| OTTER TRAWL | 3 | 0% | 4 | 0% | 3 | 0% | 23 | 0% | |
| DREDGE | 126 | 2% | 12 | 0% | 25 | 0% | 357 | 1% | |
| OTHER GEARS | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | |
| | | 4000/ | 1 | 4000/ | 1 | 1000/ | | 1000/ | |
| ALL AREAS | 5,833 | 100% | 3,831 | 100% | 5,537 | 100% | 45,815 | 100% | |
| OTTER TRAWL | 362 | 6% | 153 | 4% | 285 | 5% | 2,693 | 6% | |
| DREDGE OTHER GEARS | 5,471 0 | 94% 0% | 3,677 1 | 96% 0% | 5,251 1 | 95% 0% | 43,120 2 | 94% 0% | |
| LANDINGS - ALL AREAS | U | U /0 | ' | U /0 | ı | 0 /0 | | 0 70 | |
| Fishing Year 2003/2004 | 5,833 | | 3,831 | | 5,537 | | 45,815 | | _ |
| Fishing Year 2002/2003 | 6,298 | | 4,700 | | 3,714 | | 42,530 | | 51,734 |
| Fishing Year 2001/2002 | 4,511 | | 3,799 | | 3,906 | | 42,530 37,618 | | 47,506 |
| Fishing Year 2000/2001 | 3,645 | | 2,679 | | 2,845 | | 27,183 | | 34,309 |
| Fishing Year 1999/2000 | 2,374 | | 1,098 | | 2,193 | | 18,384 | | 22,563 |
| cg . caccc,2000 | _, | | ., | | _, | | . 0,00 . | | ,500 |

^{1.} The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Sea Scallop Stock Areas: Gulf of Maine: 511-515; South Channel: 521-522, 526

Georges Bank North: 561-562; Georges Bank South: 525

Southern New England: 537-539; New York Bight: 611-616

DE / MD / VA: 621-623, 625-627; VA / NC: 631-638

^{2.} Landings in meat weight.

^{3.} State landings for CT are estimated for the Mar. 2003 - Oct 2003 period.

^{4.} Gear data are based on vessel trip reports.

^{*} Fishing Year is March 1 through February 29.

Table 12.B. - Preliminary Scallop Landings by Major Ports and States, March - Oct 2003

| | | | | | | | | | STO | CK AREAS | S | | | | | | |
|-----------------|--------------|----------|---------|----------|---------|---------------|---------|---------------|---------|----------------|---------|----------|---------|----------|---------|------------------|------------------|
| PORT/ STATE | MAR - OCT 03 | GULF OF | MAINE | SOUTH C | HANNEL | GEORGE NOF | - | GEORGE SOL | - | SOUTHE ENGL | | NEW YOR | K BIGHT | DELM | IARVA | VIRGINIA CARC | / NORTH DLINA |
| | 1000 Lbs | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent |
| New Bedford, MA | 19,947 | 10 | 0% | 4,501 | 23% | 2,409 | 12% | 1,309 | 7% | 99 | 0% | 9,573 | 48% | 2,010 | 10% | 36 | 0% |
| Cape May, NJ | 6,038 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 2,430 | 40% | 3,581 | 59% | 26 | 0% |
| Norfolk, VA | 13,992 | 0 | 0% | 21 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 829 | 6% | 12,851 | 92% | 290 | 2% |
| | | | | | | | | 0 | | | | | | | | | |
| MAINE | 72 | 22 | 30% | 50 | 70% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| NEW HAMPSHIRE | 100 | 12 | 12% | 87 | 88% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| MASSACHUSETTS | 20,665 | 366 | 2% | 4,804 | 23% | 2,421 | 12% | 1,309 | 6% | 104 | 1% | 9,614 | 47% | 2,010 | 10% | 36 | 0% |
| RHODE ISLAND | 47 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 46 | 98% | 0 | 0% | 0 | 0% |
| CONNECTICUT | 1,250 | 0 | 0% | 0 | 0% | 28 | 2% | 0 | 0% | 0 | 0% | 1,206 | 96% | 16 | 1% | 0 | 0% |
| NEW JERSEY | 9,084 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 4,742 | 52% | 4,315 | 47% | 26 | 0% |
| VIRGINIA | 14,424 | 0 | 0% | 21 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 849 | 6% | 13,237 | 92% | 316 | 2% |
| OTHER NORTHEAST | 174 | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | 13 | 8% | 159 | 91% | 2 | 1% |
| TOTAL | 45,815 | 400 | 1% | 4,964 | 11% | 2,449 | 5% | 1,309 | 3% | 105 | 0% | 16,471 | 36% | 19,737 | 43% | 380 | 1% |

| | | | | GEAR | TYPES | | |
|-----------------|--------------|----------|---------|----------|---------|----------|---------|
| PORT/ STATE | MAR - OCT 03 | OTTER ' | TRAWL | DRE | DGE | OTHER | GEARS |
| | 1000 Lbs | 1000 Lbs | Percent | 1000 Lbs | Percent | 1000 Lbs | Percent |
| New Bedford, MA | 19,947 | 5 | 0% | 19,943 | 100% | 0 | 0% |
| Cape May, NJ | 6,038 | 755 | 13% | 5,283 | 87% | 0 | 0% |
| Norfolk, VA | 13,992 | 1,516 | 11% | 12,476 | 89% | 0 | 0% |
| | | | | | | | |
| MAINE | 72 | 0 | 0% | 70 | 98% | 2 | 2% |
| NEW HAMPSHIRE | 100 | 0 | 0% | 99 | 100% | 0 | 0% |
| MASSACHUSETTS | 20,665 | 17 | 0% | 20,648 | 100% | 0 | 0% |
| RHODE ISLAND | 47 | 1 | 0% | 46 | 99% | 0 | 0% |
| CONNECTICUT | 1,250 | 1 | 0% | 1,249 | 100% | 0 | 0% |
| NEW JERSEY | 9,084 | 825 | 9% | 8,259 | 91% | 0 | 0% |
| VIRGINIA | 14,424 | 1,788 | 12% | 12,636 | 88% | 0 | 0% |
| OTHER NORTHEAST | 174 | 62 | 36% | 112 | 64% | 0 | 0% |
| TOTAL | 45,815 | 2,693 | 6% | 43,120 | 94% | 2 | 0% |

 The three digit statistical areas defined below are for statistical and management purposes and may not be consistent with stock area delineation used for biological assessment (see the attached statistical chart).

Scallop stock areas: Gulf of Maine: 511-515, South Channel: 521-522, 526, Georges Bank North: 561-562,
Georges Bank South: 525, Southern New England: 537-539, New York Bight: 611-616,
Delmarva: 621-623, 625-627, Virginia/North Carolina: 631-638

- 2. State landings for CT are estimated for the Mar. Oct. 2003 period.
- 3. Landings in meat weight.
- 4. Gear data are based on vessel trip reports.